

# HIGHER RED MEAT INTAKES ≥3.5 SERVINGS/WEEK DO NOT INCREASE LDL CHOLESTEROL OR BLOOD PRESSURE

#### **STUDY DESIGN:**

Meta-analysis of 24 RCTs (Randomized Controlled Trials).

## **OBJECTIVE:**

To assess the effect of eating ≥0.5 servings\* of red meat/day on blood lipids, lipoproteins and blood pressure.

\*1 serving = 70 g or 2.5 oz cooked red meat.

#### **RESULTS:**

• CVD risk markers decreased in all subjects over time, with no difference in response between groups who consumed ≥0.5 or <0.5 servings of red meat/day:



- Total cholesterol
- LDL cholesterol
- Total: HDL cholesterol ratio
- Triglycerides
- Diastolic blood pressure
- Median red meat intake in the intervention diets was 2 servings/day (140 g) more than double what the average Canadian consumes.
- The highest category of red meat intake, 3 servings/day, did not increase LDL cholesterol or blood pressure.

### **STRENGTH:**

The authors note this is the first systematically searched meta-analysis of RCTs to assess the effect of ≥0.5 servings of red meat/day on blood lipids and blood pressure. Unlike observational studies, RCTs support conclusions regarding cause and effect.

## **CONCLUSION:**

Diets higher in red meat (i.e., ≥3.5 servings/week) have no adverse effect on clinically relevant CVD risk factors, compared to diets with little to no red meat.

